

INDEX TO VOLUME 81 (1995)

AUTHORS

- Adams, P.B.: see Laidig, Adams, Baxter, and Butler.
- Araya-Núñez, O., B. Ganning, and F. Bückle-Ramirez: Embryonic development, larval culture, and settling of American-pearl oyster (*Pteria sterna*, Gould) spat, 10-21.
- Barlow, G.W.: see Gillooly and Barlow.
- Barnhart, R.A.: see Busby and Barnhardt.
- Barrett, R.H.: see Kucera, Zielinski, and Barrett.
- Barrett, R.H.: see Zielinski, Kucera, and Barrett.
- Bartley, D.: see Phelps, Bartley, and Hedgecock.
- Barton, B.D.: Observations on recruitment in curl-leaf mountain-mahogany as a result of selective pruning, 85-87.
- Baxter, C.H.: see Laidig, Adams, Baxter, and Butler.
- Bleich, V.C.: see Cronin and Bleich.
- Boroski, B.B.: see Kie and Boroski.
- Bückle-Ramirez, F.: see Araya-Núñez, Ganning, and Bückle-Ramirez.
- Burton, S.F. and F. Schmalenberger: The use of Telazol for chemical restraint of black bears (*Ursus americanus*) in northern California, 29-32.
- Busby, M.S. and R.A. Barnhart: Potential food sources and feeding ecology of juvenile fall chinook salmon in California's Mattole River lagoon, 133-146.
- Butler, J.L.: see Laidig, Adams, Baxter, and Butler.
- Crabtree, R.L.: see Shiyik and Crabtree.
- Cronin, M.A. and V.C. Bleich: Mitochondrial DNA variation among populations and subspecies of mule deer in California, 45-54; errata, 170.
- Davis, J.B.: see Jensen, Petersen, Hubbart, and Davis.
- Fleming, K.J.: see Matern and Fleming.
- Fritts, T.H.: see McCoid and Fritts.
- Ganning, B.: see Araya-Núñez, Ganning, and Bückle-Ramirez.
- Garrison, B.A.: In Memoriam, David C. Zeiner, 132.
- Gillooly, J.F. and G.W. Barlow: Effects of winter floods on threespine sticklebacks in a restored stream, 155-162.
- Harvey, J.T., R.C. Helm, and G.V. Morejohn: Food habits of harbor seals inhabiting Elkhorn Slough, California, 1-9.
- Hedgecock, D.: see Phelps, Bartley, and Hedgecock.
- Helm, R.C.: see Harvey, Helm, and Morejohn.
- Hernandez, J.: see Seigel, Long, Rounds, and Hernandez.
- Hubbart, M.D.: see Jensen, Petersen, Hubbart, and Davis.
- Jensen, T., D. Petersen, M.D. Hubbart, and J.B. Davis: Use of wood duck nest boxes by swarming honey bees in the California's Central Valley, 167-170.
- Keay, J.A.: Accuracy of cementum age assignments for black bears, 113-121.
- Keay, J.A.: Black bear reproductive rates in Yosemite National Park, 122-131.
- Kie, J.G. and B.B. Boroski: Using spotlight counts to estimate mule deer population size and trends, 55-70.
- Kucera, T.E.: Recent photograph of a Sierra Nevada red fox, 43-44.
- Kucera, T.E., W.J. Zielinski, and R.H. Barrett: Current distribution of the American marten, *Martes americana*, in California, 96-103.

- Kucera, T.E.: see Zielinski, Kucera, and Barrett.
- Laidig, T.E., P.B. Adams, C.H. Baxter, and J.L. Butler: Feeding on euphausiids by *Octopus rubescens*, 77-79.
- Lea, R.N. and H.J. Walker, Jr.: Record of the bigeye trevally, *Caranx sexfasciatus*, and Mexican lookdown, *Selene brevoorti*, with notes on other carangids from California, 89-95.
- Long, D.J.: see Seigel, Long, Rounds, and Hernandez.
- Matern, S.A. and K.J. Fleming: Invasion of a third Asian goby, *Tridentiger bifasciatus*, into California, 71-76.
- McCoid, M.J. and T.H. Fritts: Female reproductive potential and winter growth of African clawed frogs (Pipidae: *Xenopus laevis*) in California, 39-42.
- Mellink, E.: Status of the muskrat in the Valle de Mexicali and Delta Del Rio Colorado, Mexico, 33-38.
- Morejohn, G.V.: see Harvey, Helm, and Morejohn.
- Petersen, D.: see Jensen, Petersen, Hubbart, and Davis.
- Phelps, A., D. Bartley, and D. Hedgecock: Electrophoretic evidence for multiple mating in tule perch, 147-154.
- Rounds, J.M.: see Seigel, Long, Rounds, and Hernandez.
- Ruiz-Campos, G.: First occurrence of the yellow bullhead, *Amieurus natalis*, in the lower Colorado River, Baja California, Mexico, 80-81.
- Schmalenberger, F.: see Burton and Schmalenberger, 29-32.
- Schulz, P.K.: Prehistoric fish remains, including thicktail chub, from the Pajaro River system, 82-84.
- Seigel, J.A., D.J. Long, J.M. Rounds, and J. Hernandez: The tiger shark, *Galeocерdo cuvier*, in coastal southern California waters, 163-166.
- Shivik, J.A. and R.L. Crabtree: Coyote activity levels in relation to presence of California gulls at Mono Lake, California, 22-28.
- Walker, H.J. Jr.: see Lea and Walker.
- Zielinski, W.J., T.E. Kucera, and R.H. Barrett: Current distribution of the fisher, *Martes pennanti*, in California, 104-112.
- Zielinski, W. J.: see Kucera, Zielinski, and Barrett.

SUBJECT

- Aging: of black bear, 113-121.
- Archaeology: prehistoric fish remains, 82-84.
- Bear, black: aging, 113-121.
- Bear, black: chemical restraint of, 29-32.
- Bear, black: reproductive rates, 122-131.
- Bees, honey: use of wood duck nest boxes, 167-170.
- Bullhead, yellow: first occurrence in the lower Colorado River, 80-81.
- Carangids: notes on others from California, 89-95.
- Chub, thicktail: remains from the Pajaro River system, 82-84.
- Coyote: activity in relation to California gulls, 22-28.
- Counts, spotlight: mule deer population and trends, 55-70.
- Creek, restored urban: effects of winter floods on threespine stickleback in, 155-162.
- Culture, larval: of American pearl-oyster, 10-21.
- Deer, mule: mitochondrial DNA variation, 45-54.

- Deer, mule: population size and trends, 55-70.
 Development, embryonic: of American pearl-oyster, 10-21.
 DNA, mitochondrial: variation in mule deer, 45-54.
 Distribution, current: American marten in California, 96-103.
 Distribution, current: fisher in California, 104-112.
 Duck, wood: honey bee use of nest boxes, 167-170.
 Errata: Cronin and Bleich, mitochondrial variation in mule deer, 170.
 Euphausiids: octopus predation, 72-76.
 Feeding ecology: of juvenile fall chinook salmon, 133-146.
 Fisher: distribution in California, 104-112.
 Fish remains: prehistoric, 82-84.
 Floods, winter: effects on threespine stickleback, 155-162.
 Food habits: of harbor seals, 1-9.
 Food sources: of juvenile fall chinook salmon, 133-146.
 Fox, Sierra Nevada red: recent photograph, 43-44.
 Frogs, African clawed: reproduction and growth in California, 39-42.
 Goby, Asian: invasion into California, 71-76.
 Growth, winter: of African clawed frogs, 39-42.
 Gulls, California: and coyote activity levels, 22-28.
 Introduction: of an Asian goby, 71-76.
 Lookdown, Mexican: range extension, 84-95.
 Mahogany, curl-leaf mountain: pruning and recruitment, 85-87.
 Marten, American: current distribution in California, 96-103.
 Mattole River lagoon: food and feeding of juvenile fall chinook salmon, 133-146.
 Memoriam: David C. Zeiner, 132.
 Muskrat: in Valle De Mexicali and Delta Del Rio, Mexico, 33-38.
 Nest boxes, wood duck: use by honey bees, 167-170.
 Octopus: feeding on euphausiids, 77-79.
 Oyster, American pearl: development and settling, 10-21.
 Population: size and trends in mule deer, 55-70.
 Range extension: bigeye trevally and Mexican lookdown, 89-95.
 Range extension: yellow bullhead, 80-81.
 Recruitment: of curl-leaf mountain mahogany, 85-87.
 Reproductive potential, female: of African clawed frogs, 39-42.
 Seals, harbor: in Elkhorn Slough, California, 1-9.
 Shark, tiger: in southern California, 163-166.
 Telazol: for chemical restraint of black bears, 29-32.
 Trevally, bigeye: range extension, 89-95.

SCIENTIFIC NAMES

- | | |
|---|--------------------------------------|
| <i>Acanthogobius flavimanus</i> , 71. | <i>Canis latrans</i> , 22-28. |
| <i>Aix sponsa</i> , 167. | <i>Caranx caballus</i> , 89, 90. |
| <i>Ameiurus natalis</i> , 80. | <i>Caranx caninus</i> , 90, 92. |
| <i>Ameiurus nebulosus</i> , 80. | <i>Caranx sexfasciatus</i> , 89-93. |
| <i>Apis mellifera</i> , 167. | <i>Carcharodon carcharias</i> , 164. |
| <i>Archoplites interruptus</i> , 83. | <i>Catostomus occidentalis</i> , 83. |
| <i>Azurina hirundo</i> , 94. | <i>Cercocarpus betuloides</i> , 86. |
| <i>Brachyrhamphus marmoratus</i> , 101. | <i>Cercocarpus ledifolius</i> , 85. |

- Cercocarpus montanus*, 86.
Chloroscombrus orqueta, 90.
Corophium spinicone, 133-144.
Cymatogaster aggregata, 147.
Decapterus scombrinus, 90.
Eogammarus confervicolus, 140.
Eucyclogobius newberryi, 74.
Galeocерdo cuvier, 164.
Gasterosteus aculeatus, 136, 155.
Gila crassicauda, 83.
Gnorimosphaeroma oregoniensis, 140.
Gulo gulo, 97, 106.
Gumaga griseus, 140.
Haemulon flaviguttatum, 94.
Hippocampus ingens, 94.
Hysterocarpus traskii, 83, 147.
Larus californicus, 22-28.
Lavinia exilicauda, 83.
Martes americana, 96, 105.
Martes americana humboldtensis, 97, 99, 100.
Martes americana sierrae, 97, 99, 100.
Martes pennanti, 97, 104.
Micrometrus minimus, 147.
Naucrates ductor, 90.
Octopus rubescens, 77-78.
Odocoileus hemionus, 45-52, 55, 85.
Odocoileus hemionus californicus, 45-52.
Odocoileus hemionus columbianus, 45, 56.
Odocoileus hemionus crooki, 45-52.
Odocoileus hemionus eremicus, 45-52.
Odocoileus hemionus fuliginatus, 45-52.
Odocoileus hemionus hemionus, 45-52.
Odocoileus hemionus inyoensis, 45-52.
Odocoileus hemionus sitkensis, 45.
Oligoplites saurus, 89.
Oncorhynchus kisutch, 144.
Oncorhynchus mykiss, 136.
Oncorhynchus tshawytscha, 133, 134.
Ondatra zibethicus, 33-38.
Orthodon microlepidotus, 83.
Otus kennicottii, 168.
Phoca vitulina, 1-9.
Pinctada mazatlanica, 10.
Pomacanthus zonipectis, 94.
Pteria sterna, 10-21.
Scorpaena mystes, 94.
Selene brevoorti, 89-93.
Selene peruviana, 90, 92.
Seriola lalandi, 90, 92.
Seriola rivoliana, 90.
Sphyrna lewini, 165.
Strix occidentalis, 101.
Sturnus vulgaris, 168.
Trachinotus paitensis, 90, 92.
Trachinotus rhodopus, 90, 92.
Trachurus symmetricus, 90, 92.
Tridentiger bifasciatus, 71-75.
Tridentiger trigonocephalus, 71-75.
Tyto alba, 168.
Uraspis secunda, 90, 92.
Ursus americanus, 29, 113, 122.
Vulpes vulpes necator, 43-44.
Xenopus laevis, 39-42.

